



Date: July 30, 2025

Patient Name: Roy

Species: Canine

Medications & Administration Schedule:

Morning medication :

8:00 AM (Morning):

- * Betacor 80 mg (Human Pharmacy): 1 tablet.
- * Administer on an empty stomach.
- * Allow 1 hour before feeding.

9:00 AM (After Feeding):

- * Epicopred 5 mg tablet (Human Pharmacy): 1 tablet.
- * Gabapentin 300 mg tablet (Human Pharmacy): 1 tablet.
- * Administer after or with food.

10:00 AM (Morning):

- * Cephalexin 750 mg tablet (Veterinary Medication): 1 tablet.

Night medications:

*** 5:00 PM (Afternoon):**

- * Gabapentin 300 mg capsule: 1 capsule.
- * Administer with food.

*** 8:00 PM (Evening):**

- * Betacor 80 mg (Human Pharmacy): 1 tablet.
- * Administer on an empty stomach.

10:00 PM (Evening):

- * Cephalexin 750 mg tablet (Veterinary Medication): 1 tablet.

11:00 PM (Night):

- * Epicopred 5 mg tablet (Human Pharmacy): 1 tablet.
- * Gabapentin 300 mg capsule: 1 capsule.
- * Administer after a small quantity of food.

Important Notes:

- Gastrointestinal Symptoms: Diarrhea and stomach gases are expected to resolve within 3-7 days post-surgery.
- Betacor (Heart Medication):
 1. DO NOT DISCONTINUE ABRUPTLY. Sudden cessation can be dangerous. Always consult with the treating veterinarian for dosage review and continuous ECG monitoring.
 2. It is best taken on an empty stomach before meals.
 3. Sedation: Betacor and Gabapentin may cause increased drowsiness and lethargy in the dog. Please note that this is a normal side effect of the medication, and there is no need to worry if it occurs.
 4. Combination Warning: Betacor and Gabapentin should NOT be administered together as this will significantly increase sedation and drowsiness in the dog.
- Feeding: It is preferable to reduce food quantities while increasing the frequency of meals. Avoid sudden large intakes of water.
- Surgical Wound Care:
 1. Keep the suture area (wound) dry at all times.
 2. Prevent licking: Ensure the dog does not lick the wound, and if there are other dogs in the house, prevent them from licking it as well.
 3. Maintain E-collar: Continuously ensure the surgical E-collar is secured.
 4. Cleaning: The wound can be regularly cleaned with a tissue using 0.9% NaCl (Normal Saline) for IV infusion.
- Bathing:
 1. You may use Betadine shampoo or Cyteal shampoo to wash the dog's body every two days, especially if urinary incontinence symptoms persist, to clean any urine residue from its body.
 2. ALWAYS re-clean the surgical wound with normal saline after bathing.
- Symptoms Requiring Concern & Veterinary Consultation:
 1. Any purulent discharge from the wound.
 2. Any opening in the wound (indicating delayed healing or bacterial infection).

If you observe any concerning symptoms, please contact the treating veterinarian and visit the hospital immediately.

- Treatment Commencement Date: Thursday, July 31, 2025
- Follow-up Consultation Date: Saturday, August 02, 2025 (Please book an appointment by phone for consultation)

Dr. Nadia Essam
Dr. Paws Animal Hospital
Phone: 01117873666 / 01008886446